

COVID-19 OAP DECISION TOOL

*In absence of specific guidance from local public health professionals this CDC based tool may be used. Specific recommendations may be modified by local authorities.

Have you been in direct* contact with a person with laboratory confirmed case of COVID-19? (*direct means person to person, NOT higher order contacts such as a 'person who met a person who met a person that was sick')

Yes

No

Are you living in the same household as, being an intimate partner of, or providing care in a non-healthcare setting for a person with symptomatic laboratory-confirmed COVID-19 infection?

No

Were you within 6 feet of the sick contact (i.e. next to someone at a dinner, meeting, interview, plane seat, town hall, etc.) for a prolonged period of time (>15 minutes) or were you exposed to respiratory secretions by being directly coughed or sneezed on?

No

Were you in the same indoor environment (office spaces, cafeteria, committee/hearing rooms, etc.) for a prolonged period of time (>15 minutes), but >6 feet away and not sneezed/coughed on?

No

All other brief contacts not categorized above (such as shaking hands, taking a photo/selfie, passing in a hallway, etc.)

Symptomatic = fever (≥ 100.4 F) and/or acute lower respiratory symptoms such as cough and shortness of breath

Yes

Yes

Yes

	High/Medium Risk	Low Risk	No Identifiable Risk
Asymptomatic People	Stay home and avoid public activities; <u>Expect quarantine</u> ; Consult public health for guidance	Self- monitor symptoms for 14 days after exposure. Personal hygiene and social distancing in accord with recommendations for all members of the community.	Personal hygiene and social distancing in accord with recommendations for all members of the community.
Symptomatic People	Immediately isolate at home; Pre-notify and consult health care provider for moderate/ severe illness; Consult public health for evaluation; <u>Expect isolation for 7 days or more.</u>	Immediately isolate at home; Pre-notify and consult health care provider for moderate/ severe illness; Consult public health for evaluation.	Immediately isolate at home; Pre-notify and consult health care provider for moderate/ severe illness; Consult public health for evaluation.